

ACETAMINOPHEN DOSING (FOR EXAMPLE, TYLENOL®)

NOT recommended for babies less than 2 months old unless recommended by a healthcare professional. Give every 4–6 hours as needed for fever or pain.

NOTE: Certain concentrations (mg/mL) of acetaminophen are being phased out.

Please be sure to check the concentration on the bottle of acetaminophen before dosing.

Weight (pounds)	Children's Liquid (160 mg/5 mL)	Children's Dissolve Tablets (160 mg tabs)
12–17 lbs	2.5 mL (80 mg)	—
18–23 lbs	3.75 mL (120 mg)	—
24–35 lbs	5 mL (160 mg)	1 tablet (160 mg)
36–47 lbs	7.5 mL (240 mg)	1.5 tablets (240 mg)
48–59 lbs	10 mL (320 mg)	2 tablets (320 mg)
60–71 lbs	12.5 mL (400 mg)	2.5 tablets (400 mg)
72–95 lbs	—	3 tablets (480 mg)

IBUPROFEN DOSING (FOR EXAMPLE, MOTRIN® AND ADVIL®)

For infants and children **OLDER** than 6 months of age. Give every 6–8 hours as needed for fever or pain.

NOTE: Check concentration on package before dosing.

Weight (pounds)	Children's Liquid (100 mg/5 mL)	Junior Tablets (100 mg)
12–17 lbs	2.5 mL (50 mg)	—
18–23 lbs	4 mL (80 mg)	—
24–35 lbs	5 mL (100 mg)	1 tablet (100 mg)
36–47 lbs	7.5 mL (150 mg)	1.5 tablets (150 mg)
48–59 lbs	10 mL (200 mg)	2 tablets (200 mg)
60–71 lbs	12.5 mL (250 mg)	2.5 tablets (250 mg)
72–95 lbs	—	3 tablets (300 mg)



Find an ER near you at **KCFastER.com**. To find a pediatric physician or answers to your health questions 24/7, call **(913) 541-KIDS (5437)**

DOSIS DE ACETAMINOFÉN (POR EJEMPLO, TYLENOL®)

NO se recomienda para bebés menores de 2 meses de edad. Dar medicamento cada 4–6 horas para aliviar síntomas de fiebre o dolor. **NOTA:** En algunas concentraciones (mg/ml) de acetaminofén las fases se separan. Asegúrese de verificar la concentración en el frasco de acetaminofén antes de administrarla.

Peso (kilos)	Jarabe líquido para niños (160 mg/5 mL)	Tabletas masticables (160 mg tabletas)
12–17 libras	2.5 mL (80 mg)	—
18–23 libras	3.75 mL (120 mg)	—
24–35 libras	5 mL (160 mg)	1 tableta (160 mg)
36–47 libras	7.5 mL (240 mg)	1.5 tabletas (240 mg)
48–59 libras	10 mL (320 mg)	2 tabletas (320 mg)
60–71 libras	12.5 mL (400 mg)	2.5 tabletas (400 mg)
72–95 libras	—	3 tabletas (480 mg)

DOSIS DE IBUPROFENO (POR EJEMPLO, MOTRIN® Y ADVIL®)

Para niños **mayores** de 6 meses. Dar medicamento cada 6–8 horas para aliviar síntomas de fiebre o dolor.

NOTA: Verifique la concentración en el paquete antes de la dosificación.

Peso (kilos)	Jarabe líquido para niños (100 mg/5 mL)	Tabletas para niños (100 mg)
12–17 libras	2.5 mL (50 mg)	—
18–23 libras	4 mL (80 mg)	—
24–35 libras	5 mL (100 mg)	1 tableta (100 mg)
36–47 libras	7.5 mL (150 mg)	1.5 tabletas (150 mg)
48–59 libras	10 mL (200 mg)	2 tabletas (200 mg)
60–71 libras	12.5 mL (250 mg)	2.5 tabletas (250 mg)
72–95 libras	—	3 tabletas (300 mg)



Find an ER near you at **KCFastER.com**. To find a pediatric physician or answers to your health questions 24/7, call **(913) 541-KIDS (5437)**